



# ORIGINAL ARTICLE

## Evaluation of Relation between Psychiatric Capital and Prevalence of Postpartum Anxiety and Depression in Different health Centers of Birjand

Ali Nadjafi-Semnani<sup>1</sup>, Yahya Mohammadi<sup>2</sup>, Fatemeh Nadjafi-Semnani<sup>3</sup>  
 Mohammad Nadjafi-Semnani<sup>4</sup>, Asieh Shahriyari<sup>5</sup>, Nahid Ghanbarzadeh<sup>6</sup> ✉

<sup>1</sup> Medical Student, Tehran University of Medical Sciences, Tehran, Iran

<sup>2</sup> EDC, Birjand University of Medical Sciences, Birjand, Iran and PhD Student of Curriculum Development, Allameh Tabataba'i University, Tehran, Iran

<sup>3</sup> Medical Student, Tehran University of Medical Sciences, Tehran, Iran

<sup>4</sup> Department of Urology, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>5</sup> MA of psychology and psychiatrics, Shahid Rahimi hospital, Birjand, Iran

<sup>6</sup> Department of Gynecology and Obstetrics, Birjand University of Medical Sciences, Birjand, Iran

Received: September 9, 2017

Revised: December 27, 2017

Accepted: May 21, 2018

### Abstract

**Introduction:** Postpartum depression and anxiety is an important and disabling disorder that, if not diagnosed and treated in a timely manner, could worsen and develop into a chronic condition. Different factors play a role in this disorder, including psychiatric capitals. This study plans to investigate the relation between psychiatric capitals and prevalence of postpartum depression and anxiety in females referring to different medical centers of Birjand.

**Methods:** This was a descriptive correlational study. Women who had a history of delivery 6 to 8 weeks prior to the study were enrolled. Of all available medical centers, 5 were chosen using randomized-clustered method and 162 patients were picked. These patients were asked to fill Edinburgh's postnatal depression scale (EPDS), Zung's self-rating anxiety scale (SAS) and Lathan's psychological capital questionnaires. Data were gathered and analyzed using SPSS version 18 and Pearson correlation coefficient tests.

**Results:** Mean age of participants was 32.25±4.44. There was a significant reverse correlation between psychiatric capitals and depression and anxiety of postpartum (p-value <0.05).

**Conclusions:** This study shows that psychiatric capital and its elements could predict psychological well-being accurately and reliably. With this in mind, medical centers supporting pregnant women should hold different related workshops aiming to improve psychiatric capital of their clients.

**Key Words:** Psychiatric Capital; Anxiety; Depression; Delivery

©2018 Journal of Surgery and  
 Trauma  
 Tel: +985632381203  
 Fax: +985632440488  
 Po Bax 97175-379  
 Email: jsurgery@bums.ac.ir



✉ Correspondence to:

Nahid Ghanbarzadeh, MD, Department of Gynecology and  
 Obstetrics, Birjand University of Medical Sciences, Birjand,  
 Iran;  
 Telephone Number: +9809155610029  
 Email Address: nghanbarzaedh@gmail.com

### Introduction

Depression is one of the most common  
 psychiatric disorders and a common denominator

of human life, and is seen in almost all countries and cultures. Various studies show that women are depressed 2 times more than men. Stress factors such as childbirth, menstruation, learned helplessness, hormonal differences, inadequate social skills, and the difference in psychosocial stresses in men and women are involved (1). During pregnancy and after delivery, very significant psychological and physiological changes occur that sometimes lead to pathological changes and cause psychiatric disorders. Postpartum mental disorders, including postpartum depression, create problems for the mother, infant and other family members, and this may affect the level of attachment of the mother and the baby and other family relationships, and even a threat to safety and health of mother, baby and other children (2-5).

Anxiety is another mental disorder that occurs in pregnant women and after childbirth. In many women, the birth of a child is the cause of anxiety that triggers the onset or recurrence of chronic depression disorders (6). In fact, the postnatal period is a transitional period in which the mother faces new patterns and relationships that requires adaption (7).

Among the variables that affect the mental health of mothers during pregnancy and delivery is psychological capital, which is one of the indicators of positivism, which is characterized by individual belief in one's ability to succeed, perseverance in pursuing goals, creation of positive attributes about oneself and tolerance of problems (8). Also, being able to have psychological capital enables individuals to be less tense, in addition to better coping with stressful situations, and to have high potential in order to face problems. (9). Seligman believes that psychological capital embodies the positive aspects of human life. He believes that human and social capital is obvious, easily observable and can be easily measured and controlled. While psychological capital is more potent, and is difficult to measure and develop (10, 11).

Recent studies at the University of Michigan suggest that there is a relationship between the symptoms of maternal depression and hormonal neurological changes and adjustment disorders in various aspects of sleep and nutrition and mood and dependence in the newborn (12). Depressed mothers are less responsive to the infant, and they have complex problems with the baby (13), which seriously threatens the emotional-cognitive development of the infant and may lead to negligence and misbehavior in the baby's care (14, 15). The cause of this disease is not yet known;

however, biological factors such as sudden loss of estrogen concentration after delivery, decreased progesterone, increased cortisol urinary excretion, anti-thyroid antibodies, prolactin, oxytocin and beta-endorphins (16 and 17) are involved. Therefore, timely diagnosis and differentiation of this disorder from postpartum depression is one of the tasks of the treatment group. The medical team (including general practitioner, midwife, and family planning and vaccination experts) should be able to identify those who are prone to postnatal psychiatric disorders (especially depression and postpartum anxiety), and guide them regarding the care and support for this period (18). Treatment with serotonin reuptake-inhibitors, psychotherapy (sometimes as a treatment option and sometimes to increase the acceptance of drug therapy by the patient), and in severe cases, hospitalization are among the treatments of this disorder (19). Therefore, the symptoms of depression during pregnancy or previous history of depression and the presence of risk factors necessitate careful follow-up (20). This study was conducted to investigate the relationship between psychological capital and postpartum depression and anxiety in women in Birjand.

## Methods

This was a descriptive correlational study. The statistical population consisted of all mothers who referred to health centers during the first 6 to 8 weeks postpartum. From the health centers of the city, 5 centers were selected randomly and then 162 people (according to Morgan table) were selected by available sampling. The criteria for inclusion into the study were all women who referred to health centers within 6-8 weeks postpartum and were willing to participate in the study. In order to collect data, the Edinburgh, Zang, and Luthanes questionnaires were used. The Standard Depression Inventory of Edinburgh has 10 questions related to depression, anxiety, feelings of guilt and suicidal thoughts, and has been set up by Cox et al. (1987) and has been used repeatedly to investigate the diagnosis of postpartum depression. The questionnaire is designed to make it possible to diagnose depression from 6 weeks postpartum. The Edinburgh Score is between 0 and 30, and score 12 or more is considered as postpartum depression. Questions 1, 2, 4 are scored from 0 to 3 and questions 3, 5, 6, 7, 8, 9, 10 are scored from 3 to 0 (19). The Zang Anxiety Inventory was designed by Zang in 1970. The Zang Anxiety Scale is designed to measure anxiety and has 20 questions with 4

choices: never, occasionally, most often and almost always. The questions in this questionnaire are based on physical-emotional syndrome. Since some questions are scored inverted, fewer references can give the same answer to all options. Also, a number of questions on this scale emphasize the positive symptoms of anxiety and a number of others emphasize on the negative symptoms. Those with less anxiety have a lower score and those who are more anxious get higher scores on this scale (6). Also, the Luthanese psychological capital inventory questionnaire consists of 24 questions and 4 subscales of hope, tangibility, optimism and self-efficacy in which each subscale has 6 sentences; questions 1-6 are related to the self-efficacy subscale, questions 7-12 are related to the hope subscale, questions 13 to 18 are related to the resilience subscale and questions 19-24 are related to the optimistic subscale (22). The questionnaires were validated by three faculty members. To determine the reliability of the questionnaires, the Cronbach's alpha method was

mean anxiety score was  $39.25 \pm 9$ , the mean score of depression was  $18 \pm 8$  in the studied women. The Kolmogorov Smirnov test was used to determine the normality of the data. All variables had normal distribution due to their significance level of more than 0.05.

The results also showed that there is a significant negative relationship between psychological capital components with postpartum depression and can predict the level of depression based on the components of psychological capital ( $p < 0.05$ ). Also, there is a significant negative relationship between the components of psychological capital and anxiety and the level of anxiety can be predicted based on the components of psychological capital ( $p < 0.05$ ) (Table 1).

## Discussion

The main objective of this study was to determine the relationship between psychological capital and depression and anxiety in women after

**Table 1: Relationship between dimensions of psychological capital with postpartum depression and anxiety**

Optimism (N=162)	Resilience (N=162)	Hope (N=162)	Self-efficacy (N=162)	Psychological capital (N=162)	Variable
R=-0.520 P=0.0001	R=-0.474 P=0.0001	R=-0.484 P=0.0001	R=-0.132 P=0.02	R=-0.508 P=0.0001	Postpartum depression
R=-0.318 P=0.0001	R=-0.279 P=0.0001	R=-0.193 P=0.015	R=-0.068 P=0.41	R=-0.242 P=0.002	Anxiety

used and the Cronbach's alpha coefficient was 85%, 79% and 81%, respectively. Data analysis was performed using SPSS software version 18 and descriptive statistics including mean, standard deviation and inferential statistics including Pearson correlation coefficient at the significance level of 0.05 were performed.

## Results

The mean age of women was  $32.25 \pm 4.44$ . The highest number of respondents was between the ages of 27 to 31 (41.36%) and the lowest number of respondents were from 17 to 21 (12.34%) and based on the degree of education, 27.77% had diploma, 50% had bachelor's degree and 21.6% had master's degree.

Also, the results showed that the average score of women's hope was  $21.5 \pm 6$ , the mean score of optimism was  $20.9 \pm 5$ , the average score of resilience was  $25 \pm 6$ , self-efficacy was  $23 \pm 9$  and the total psychological capital was  $90 \pm 23$ , the

childbirth. Childbirth is one of the most influential changes in women's lives. This change is an unforgettable experience not only for women, but due to its importance, it is very effective in shaping the psychological and physical effects of women (21). The results of this study showed that psychological capital and its components have a negative and significant relationship with depression in postpartum women. With the increase in psychological capital, women's depression reduces. These findings are consistent with the results of Hughes et al. (22). The psychological capital is a function of social capital, and the quantity and quality of interactions in social relations determine social capital. On the other hand, social relationships affect the size of psychological capital. Therefore, changes in the quantity and quality of these social relationships can affect the performance of a person. Adler et al. (23) also showed that the fluctuations and changes in the status of psychological capital at the individual and general levels are correlated with the change in the amount of social capital. Thus,

social capital is associated with a more favorable situation of individual and general health, higher levels of optimism, increased positive health behaviors, increased resilience to problems, and reduced mental disorders. People with a high psychological capital have the ability to interact more socially. So, what connects people to each other, and leads to repetition and continuity of interactions, is psychological capital. It seems that people with a higher psychological capital can have more continuous social interactions and better social life.

Further findings showed that there is a significant negative relationship between psychological capital and anxiety in postpartum women. Therefore, psychological capital is directly related to happiness and well-being of individuals and psychological well-being prevents anxiety in difficult conditions. This conclusion is not consistent with the investigations of Decker (24), Saadati et al. (25). Therefore, the anxiety during pregnancy can affect mother's relationship with the infant and affect the mental health of the children in the next evolutionary period, and the children will face mental health problems in adulthood.

One of the positive aspects of our study is that our studied women are from different social, economic and cultural classes of the society, and therefore a good representative of the society. Of course, the study method is cross-sectional and for mothers immediately after delivery. Therefore, it is recommended that further studies be carried out in a cohort manner. The present study can be used in midwifery and psychological services to promote maternal and child health especially in reducing the causes of depression and anxiety and stress in mothers.

## Conclusions

Since psychological capital had a significant correlation with depression and anxiety variables in postpartum women, it is suggested that the grounds for the increase of this capital, both at the family level and at the community level, are prepared. The existence of a significant amount of psychological capital facilitates social activities, so that in times of crisis, it can be used to solve problems as the main source of solving problems such as depression, and the current study confirms this issue.

## Acknowledgements

Researchers are grateful to women who have provided the ground for this research with their contributions, and wish them all the best. This

study was approved ethically (ethical code: 1396.328 Ir.bums.REC)

## Author's contribution

Ghanbarzdeh N, Shahriyari A and Nadjafi-Semnani A designed the study. Shahriyari A helped with data gathering and coordination with different medical centers. Final data analysis was performed by Mohammadi Y and Nadjafi-Semnani A. Manuscript was prepared by Nadjafi-Semnani M, Mohammadi Y and Nadjafi-Semnani F, Nadjafi-Semnani A and Ghanbarzdeh N. All authors read and approved the final manuscript.

## Funding

This study has not been funded by any natural/legal person.

## Conflict of interest

There is no conflict of interest of authors for this study.

## References

1. Sadock B, Sadock V. Kaplan and Sadock's synopsis of psychiatry-behavioral sciences/clinical psychiatry. 10<sup>th</sup> ed. Philadelphia: Lippincott Williams and Wilkins; 2007.
2. Janati Y, Khaki N. [Psychiatry in midwifery]. 1<sup>st</sup> ed. Tehran: Jamee Negar; 2005. [Persian]
3. Doulatian M, Maziyar P, Alavi Majd Hamid, Yazdjerdi M. [The relationship between mode of delivery and postpartum depression]. *J Reprod Infertility*. 2006;7(3):260-8. [Persian]
4. Hosseini H, Naghibi AH, Khademloo M. [Postpartum depression and its relationship with some related factors]. *J Babol Univ Med Sci*. 2008;10(2):76-81. [Persian]
5. Perfetti J, Clark R, Fillmore CM. Postpartum depression: identification, screening and treatment. *WMJ*. 2004;103(6):56-63.
6. Robertson E, Grace S, Wallington T, Stewart DE. Antenatal risk factors for postpartum depression: a synthesis of recent literature. *Gen Hosp Psychiatry*. 2004 Jul-Aug;26(4):289-95. DOI: 10.1016/j.genhosppsy.2004.02.006
7. Stocky A, Lynch J. Acute psychiatric disturbance in pregnancy and the puerperium. *Baillieres Best Pract Res Clin Obstet Gynaecol*. 2000 Feb;14(1):73-87.
8. Luthans F, Luthans KW, Luthans BC. Positive psychological capital: going behind human and

- social capital. *Bus horiz.* 2004 Jan-Feb;47(1):45-50.
- 9- Robbins S, Millet B, Caccioppe R, Waters- Marsh T. *Organizational behavior.* 2<sup>nd</sup> ed. Australia: Prentice Hall; 1998.
  10. Seligman ME, Csikszentmihalyi M. *Positive psychology: An introduction.* *Am Psychol.* 2000 Jan;55(1):5-14.
  11. Goldsmith AH, Veum JR, Darity Jr W. Unemployment, joblessness, psychological wellbeing and self-esteem: Theory and evidence. *J Socio Econ.* 1997; 26(2):133-58. DOI: 10.1016/S1053-5357(97)90030-5.
  12. Marcus SM. Depression during pregnancy: rates, risks and consequences--Motherisk Update 2008. *Can J Clin Pharmacol.* 2009 Winter;16(1):e15-22.
  13. Posmontier B. Sleep quality in women with and without postpartum depression. *J Obstet Gynecol Neonatal Nurs.* 2008 Nov-Dec;37(6):722-35; quiz 735-7. doi: 10.1111/j.1552-6909.2008.00298.x.
  14. Vesga-Lopez O, Blanco C, Keyes K, Olfson M, Grant BF, Hasin DS. Psychiatric disorders in pregnant and postpartum women in the United States. *Arch Gen Psychiatry.* 2008 Jul;65(7):805-15. doi: 10.1001/archpsyc.65.7.805.
  15. Kendall-Tackett K. A new paradigm for depression in new mothers: the central role of inflammation and how breastfeeding and anti inflammatory treatments protect maternal health. *Int Breastfeed J.* 2007; 2: 6. DOI: 10.1186/1746-4358-2-6.
  16. McCoy SJ, Beal JM, Shipman SB, Payton ME, Watson GH. Risk factors for postpartum depression: a retrospective investigation at 4- weeks postnatal and a review of the literature. *J Am Osteopath Assoc.* 2006 Apr;106(4):193-8.
  17. Nakku JEM, Nakasi G, Mirembe F. Postpartum major depression at six weeks in primary health care: prevalence and associated factors. *Afr Health Sci.* 2006 Dec;6(4):207-14.
  18. Rasouli P, Shobeiri F, Cheraghi F, Rasouli R, Ghanbari V. Study of the relationship of anxiety and depression in third trimester pregnancy on growth index of neonates and preterm delivery. *Iran J Pediatr Nurs.* 2016;2(4):1-9. [Persian]
  19. Salary P, Banafshe E, Hebrani P, Jabbari Nooghabi H. [On the relationship between maternal fatigue and postpartum depression]. *J Fundam Ment Health.* 2010; 11(4): 302-11. [Persian]
  20. Hosseini SH, Naghibi AA, Khademloo M. [Postpartum depression and its relationship with some related factors]. *J Babol Univ Med Sci.* 2008 Jun-Jul; 10(2): 76-81. [Persian]
  21. Khodabakhshi Koolae A, Heidari S, Khoshkonesh A, Heidari M. [Relationship between Spiritual Intelligence and Resilience to Stress in Preference of Delivery Method in Pregnant Women]. *Iran J Obstet Gynecol Infertil.* 2013;16(58):8-15. [Persian]
  22. Hughes LW. A correlational study of the relationship between sense of Humor and positive psychological capacities. *Economics & Business Journal: Inquiries & Perspectives.* 2008;1(1):46-55.
  23. Adler PS, Kwon SW. Social capital: Prospects for a new concept. *Acad Manage Rev.* 2002 Jan;27(1):17-40. DOI: 10.2307/4134367
  24. Dekker WH, Rotondo DM. Relationships among gender, Type of humor, and perceived leader effectiveness. *Journal of Managerial Issues.* 2001 Winter;13(4):450-65.
  25. Sa'adati M; Abbas Zadeh M; Niazi M; Montazery Khosh H. [Sociological study of the relationship between social capital and depression (Case study: married women of Tabriz city)]. *Quarterly Journal of Women's Studies Sociological and psychological.* 2015;13(1):89-110. DOI: 10.22051/jwsp.2015.1939. [Persian]