

ORIGINAL ARTICLE

Accident-related mortality in Southern Khorasan Province in 2010

Toba Kazemi¹, Ahmad Amouzesi²✉, Nahid Borna³, Gholam Reza Sharifzadeh⁴,
Zahra Amouzesi⁵

¹Professor, Atherosclerosis and Coronary Artery Research Centre, Birjand University of Medical Sciences, Birjand, Iran;

²Assistant professor, Atherosclerosis and Coronary Artery Research Centre, Birjand University of Medical Sciences, Birjand, Iran;

³member of the Student Research Committee, MD, Birjand University of Medical Sciences, Birjand, Iran;

⁴Assistant Professor, Social Determinants of Health Research Center, Birjand University of Medical Sciences, Birjand, Iran;

⁵Instructor, Faculty of Nursing and Midwifery, Birjand University of Medical Sciences, Birjand, Iran.

Received: February 23, 2015 Revised: June 24, 2015 Accepted: June 24, 2015

Abstract

Introduction: Causes of death vary globally, and they depend regionally on lifestyle and socio-cultural factors. The purpose of this study was to determine accident-related mortality in Southern Khorasan Province, Iran, in the year 2010.

Methods: This descriptive-analytical study involved all the death cases in the year 2010 in Southern Khorasan. The data were collected from the death registration system, and were classified and analyzed according to the codes of the 10th revision of the International Classification of Deaths (ICD).

Results: From among the 3,792 deaths in 2010, n=633 (16.7%) were accident-related. Of accident-related deaths, 73.6% (n=450) of the deceased persons were male. The highest rate of mortality belonged to Summer (July 23 to Aug 22) (n=203; 32.1%). The median age in accident-related deaths was 36.6±21.9 years with the minimum age of 5 days and maximum age of 92 years. The leading cause of mortality was traffic accidents (52.1%) and violence (18%). The median age of the deceased women in accidents was significantly higher than men (P<0.048) and in rural persons more than urban people (P=0.02), with Surgical complications as the least frequent cause of death.

Conclusions: Since road traffic accidents rank first in accident-related mortality, it is recommended to have measures to improve road safety as well as quality and safety of vehicles, and to establish strict rules for those who disobey traffic rules.

Key Words: mortality; intentional and unintentional injuries; accidents

Introduction

Types of deaths and diseases have been changing across the globe in recent years where accident deaths have turned into a major concern [1]. Injuries are considered as causes of fatality and inability worldwide. They account for about 16% of global bar of diseases and the cause of death for

people below 60 years [2]. The indirect estimation of World Health Organization (WHO) and the Global Burden of Diseases (GBD) indicate that unintentional accidents result in 3.9 million deaths worldwide of which 90% occur in low- and middle-income countries. The most common causes of accident mortality include road traffic accidents, drowning, poisoning, and burning [3]. The main

© 2014 Journal of Surgery and Trauma; Birjand University of Medical Sciences Journal Office, Ghaffari Ave., Birjand, I.R. Iran
Tel: +985632443041 (5533)
Fax: +985632440488
Po Bax 97175-379
Email: jsurgery@bums.ac.ir



✉ Correspondence to:

Ahmad Amouzesi; Assistant professor, Atherosclerosis and Coronary Artery Research Centre, Birjand University of Medical Sciences, Birjand, Iran.
Telephone Number: +05632440488
Email Address: amouzesiahmad@bums.ac.ir

causes of death among the Iranian population consist of cardiovascular diseases, motor vehicle accidents, and intentional and unintentional injuries, respectively [4]. The majority of the injuries can be prevented where their prevention is a priority in developed countries [1]. Given the importance of this issue, therefore, this study was conducted to determine the causes of Accident-related mortality in Southern Khorasan Province, in the east of Iran, in the year 2010.

Methods

This descriptive-analytical study included all the deceased persons in Southern Khorasan in 2010 according to the Iranian death registration system. First, all the death cases were collected using the death registration system form from all the health houses, health-care centers, hospitals, Legal Medicine Organization, and cemeteries. They were examined in the committee for death registration. The incomplete forms were referred back to the reporting place and later collected. The causes of death were coded according to the 10th revision of ICD and then analyzed in SPSS using one-way Anova and independent t-test at the significant level of <0.05.

Results

In this study, from among the 3,792 deaths in 2010, n=633 (16.7%) resulted from accidents in the province. Of accident deaths, 73.6% (n=450) of the deceased persons were male. The highest death rate occurred in Summer (July 23 to Aug 22) (n=203; 32.1%). Frequency distribution of accident deaths is provided in Table 1 by age, residence, and the season in which the deaths occurred. The median age in this type of mortality was 36.6±21.9 years with the minimum age of 5 days and maximum age of 92 years.

The leading cause of death was road traffic accidents (52.1%), followed by violence (18%) (Table 2). The median age of deceased women in accidents was significantly higher than men (P<0.048) and in rural persons more than urban people (P=0.02), with Surgical complications as the least frequent cause of death (Table 3).

Table 1: Frequency distribution of accident deaths by age, residence, and season

| Variable | Frequency | Percent | |
|-------------|-----------|---------|------|
| Season | Spring | 174 | 27.5 |
| | Summer | 203 | 32.1 |
| | Autumn | 114 | 18 |
| | Winter | 121 | 22.2 |
| Residence | Urban | 241 | 38.1 |
| | Rural | 392 | 61.9 |
| Age (years) | < 20 | 140 | 22.2 |
| | 20-44 | 290 | 45.8 |
| | 45-65 | 119 | 18.8 |
| | > 65 | 84 | 13.2 |

Table 2: Frequency distribution of Cause of death due to accident

| Cause of death | Frequency | Percent |
|--|-----------|---------|
| Traffic accidents | 330 | 52.1 |
| Violence by others | 114 | 18 |
| Burning | 56 | 8.8 |
| Falling | 21 | 3.3 |
| Unintentional accidents | 20 | 3.2 |
| Suicide | 17 | 2.7 |
| Accidental poisoning by drugs and medicine | 14 | 2.2 |
| Surgical complications | 13 | 2.1 |
| Other reasons | 48 | 7.6 |
| Total | 633 | 100 |

Table 3. Comparison of median age in accident deaths by Sex, residence, and cause of death

| Variable and sub-groups | Age | | |
|-------------------------|-------------------------|------|-------------------------------------|
| | Frequency | X±SD | P value |
| Sex | Male | 450 | 39.6±26.6 |
| | Female | 163 | 35.5±19.9 |
| Residence | Urban | 238 | 34.1±20.5 |
| | Rural | 375 | 38.3±22.7 |
| Cause of death | Unintentional accidents | 496 | 37.9±23.8 |
| | Violence by others | 100 | 31.9±9.4 |
| | Suicide | 17 | 26.9±8.9 |
| | | | unintentional accidents or violence |
| | | | 0.03 |
| | | | Unintentional accidents or suicide |
| | | | p<0.001 |

Discussion

In the present study, three common causes of accident-related mortality were Traffic accidents, Violence by others, and Burning. According to Saadat (2014), the leading causes of mortality among Iranians include cardiovascular diseases, motor vehicle accidents, cancers, and intentional and unintentional injuries, respectively. Life years lost root respectively from cardiovascular diseases, motor vehicle accidents, cancers, and intentional and unintentional injuries [4]. These findings are similar to those of other countries.

In our study, the most common cause of death was accidents. According to WHO estimates, the most frequent causes of unintentional accident deaths include road traffic accidents, falling, drowning, poisoning, and burning [3]. WHO predicts that road traffic accidents will be on the rise by 2020 except for high-income countries [5]. Road traffic accidents are the second commonest cause of death in Iran, while they are the ninth at the global scale and seventh in the United States [6].

In our study, men were the major victims in accident deaths. In Khuzestan Province, in the southeast of Iran, similar findings were observed in the same year (82% male; 18% female).

In Southern Khorasan in 2010, the majority of deaths occurred in the age group of 20-44 years. In Ghorbani Birghani (2012), the highest rate of mortality belonged to the age group of 15-44 years [6] which is somehow similar to our findings. This highlights the necessity on the part of authorities for immediate action in identifying and preventing the causes. In our study, accident deaths occurred more during the summer. Chen (2013) showed that accident deaths in Canada followed an obvious seasonal pattern in which men were the main victims. The reason mentioned for the seasonal pattern can be summer vacations and increased travels [2].

In the current study, the leading cause of death after road traffic accidents (52.1%) was violence (18%). In 2012, mortality rate due to violence was 10% at the global scale and 7% in the eastern Mediterranean region [7]. In the same year, violence mortality rates in men and women were 1.6% and 5.2% respectively [8], which are lower than the rates in Iran. When their risk factors are known, the majority of injury and violence causes of death can be prevented. Nonetheless, violence and injury-related risk factors are complicated and multi-factored. Interventions, on the other hand, can both reduce mortality and have economic

implications because much of the medical, social, and legal expenses can be saved [7].

Burning mortality in the present study counted to 8.27 per 100,000 persons. In Amiralavi (2013), burning mortality was reported as 8.7. In the developed countries, however, the rate is 2.1 per 100,000. Mortality risk factors vary across countries and depend regionally on lifestyle and socio-cultural factors [9]. In Amiralavi's study, risk factors of burning mortality are indicated by adulthood, illiteracy, unemployment, residence in rural areas, and low socio-economic status along with burning percents. The difference between urban and rural areas resides in access to medical and emergency services that can affect mortality rate [10]. Another probable reason for the higher rate of mortality in rural areas is the greater contact of rural residents with environmental injury factors than the urban residents.

In the current study, mortality arising from falling was 33%, while the rate is 5% in the Mediterranean region and 14% worldwide [7]. In this study, poisoning rate was 22%; this rate was 4% in the Mediterranean region and 14% worldwide [7]. Global poisoning mortality rates in the year 2010 in men and women (for 14-49 year-olds) were reported as 9% and 1.2% [9]. Generally speaking, falling and poisoning mortality in the current study was less than other regions. The reason can be the fewer number of the elderly population in Iran than other countries because this population is more prone to falling than other age groups. Unintentional poisoning usually occurs as a result of chemicals in houses, medicine, and plants. This can be due to the more limited access to chemicals in Southern Khorasan province because of the limited number of factories and greenhouses.

Suicide was the least frequent cause of death in the current study (2.5 per 100,000). It is the 13th cause of death across the world [11]. In a systematic review of suicide in Iran, suicide had a frequency 9.4 per 100,000 [12]. The reason for the lower rate of suicide in our study than the mean rate in Iran can be the cultural status of the region in terms of social literacy and religious beliefs.

Conclusions

To prevent road traffic deaths, it is recommended to have measures to improve road safety as well as quality and safety of vehicles, and to establish strict rules for those who disobey traffic rules. In addition, elevated coordination between different centers such as police stations, fire department, and emergency medical services

can provide improved and faster services to victims.

Acknowledgements

This paper draws upon the dissertation thesis of Nahid Borna. The authors deeply appreciate the Health Division of Birjand University of Medical Sciences and Ms Kafayi that made great contributions in data collection.

References

1. Akbari ME, Naghavi M, Soori H. Epidemiology of deaths from injuries in the Islamic Republic of Iran. *East Mediterr Health J.* 2006;12(3-4):382-90.
2. Chen Y, Mo F, Yi QL, Jiang, Y, Mao Y. Unintentional injury mortality and external causes in Canada from 2001 to 2007. *Chronic Dis Inj Can.* 2013;33(2):95-102.
3. Jagnoor J1, Suraweera W, Keay L, Ivers RQ, Thakur J, Jha P; Million Death Study Collaborators. Unintentional injury mortality in India, 2005: Nationally representative mortality survey of 1.1 million homes. *BMC Public Health.* 2012;12:487.
4. Saadat S, Yousefifard M, Asady H, Moghadas Jafari A, Fayaz M, Hosseini M. The most important causes of death in Iranian population; a retrospective cohort study. *Emergency.* 2015;3(1):16-21.
5. Bahadori Monfared A, Soori H, Mehrabi Y, Rahmati Roudsar M, Esmaili A, Salehi M, et al. A model for prediction of on the rate of mortality due to road traffic accidents in Iran. *Journal of the faculty of medicine, Shaheed Beheshti University of Medical Sciences and Health Services.* 2013;36(5):7-11. [Persian]
6. Ghorbani Birgani AR, Hakim AS, Zare K. Epidemiologic study of fatal traffic accidents in khuzestan province in 2010. *Quarterly Journal of Rescue & Relief.* 2012;4(2):28-35. [Persian]
7. Bachani AM, Zhang XJ, Allen KA, Hyder AA. Injuries and violence in the Eastern Mediterranean Region: a review of the health, economic and social burden. *East Mediterr Health J.* 2014;20(10):643-52.
8. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet.* 2012;380(9859):2095-128.
9. Alavi SA, Mobayen MR, Tolouei M, Noursalehi I, Gholipour A, Gholamalipour N, et al. Epidemiology and outcome of burn injuries in burn patients. *Journal of Qom University of Medical Sciences.* 2013;7(5):35-41. [Persian]
10. Burrows S, Auger N, Gamache P, Hamel D. Leading Causes of Unintentional Injury and Suicide Mortality in Canadian Adults Across the Urban-Rural Continuum. *Public Health Rep.* 2013;128(6):443-53.
11. Astaraki P, Keikhavani S, Mansourian M, Bashiri S, Qorbani M, Rezapoor A, et al. A Comparative Study of the Causes and Methods of Suicide Lead to Death Referred to Ilam Legal Medicine Center in 2004-2009. *Scientific Journal of Forensic Medicine.* 2014;19(1-4):1-8. [Persian]
12. Ghoreishi SA, Mousavinasab N. Systematic review of researches on suicide and suicide attempt in Iran. *Iranian Journal of Psychiatry and Clinical Psychology.* 2008;14(2):115-21. [Persian]