

LETTER TO EDITOR

Risks of Ghaderi's Phenomenon

Reza Ghaderi¹ 

¹Professor, Department of Dermatology, Faculty of Medicine, Birjand University of Medical Science, Birjand, Iran

Received: June 20, 2019 Revised: July 30, 2019 Accepted: August 04, 2019

Dear Editor

"Beauty has so many forms, and I think the most beautiful thing is confidence and loving yourself".
Kieszka

Beauty has classically been considered amongst the definitive values, goodness, justice, integrity, and loyalty. Beauty, cutaneous body image, quality of life, self-esteem, and self-confidence are intertwined together (1-2). Life quality consists of a feeling of joyance and satisfaction with life. However, life quality, self-esteem, and self-confidence in patients with dermatoses have not adequately been attended to so far. As regards, dermatoses affect general health, well-being, and social adaptation of the individual (3-5). In a study, the majority of the female patients who were candidates for Bariatric Surgery declared high body image concerns (6). Modern cognitive approaches to Body Dysmorphic Disorder suppose that appearance-related intrusive cognitions and their functional outcomes determine the disorder (1).

The Body Dysmorphic Disorder is a common debilitating disease worldwide. Patients manifest a disabling preoccupation with a few perceived defects, frequently well-marked by poor insight or delusional thoughts (2-7). According to the results of a study, rhinoplasty candidates reported high body image concern (8). Accurate interviewing of the patients who are candidates for rhinoplasty is very important to evaluate their level of body image concern in order to diagnose any available disease and prescribe required treatments (8).

More and more people worldwide are using hyaluronic acid injections for skin rejuvenation and better cutaneous body image and self-esteem (5- 7). Cosmetic injectable treatments are progressively common. However, a few studies have investigated modifications in psychosocial functioning following these procedures (9). Some studies were performed to evaluate changes in cutaneous body image, self-esteem, and life quality following cosmetic injectable procedures with neuromodulators and soft tissue fillers (9). Cosmetic injectable treatments associated with an increase in body image satisfaction. Further investigation using modern cosmetic surgery-specific devices can supply new insights into the psychosocial advantages of cosmetic injectable treatments (9). Millions of cosmetic and surgical procedures of the face are administrated each year; however, objective clinical criteria that assess surgical procedures, such as complication rates, have limited efficiency when applied to cosmetic procedures. While there might be subjective improvements in cutaneous body image, it is very important to assess if these procedures have an effect on patients in other domains, such as psychosocial functioning (10).

Most studies showed moderate improvements in psychosocial functioning, including a self-body image, self-esteem, and life quality. Unluckily, the global quality of evidence is limited due to a lack of control groups, short follow-up periods, or loss to follow-up (10). The current evidence shows that a number of psychosocial realms can improve

©2019 Journal of Surgery and
Trauma

Tel: +985632381203

Fax: +985632440488

Po Bax 97175-379

Email: jsurgery@bums.ac.ir

✉ Correspondence to:

Reza Ghaderi, Professor, Department of Dermatology, Faculty of Medicine, Birjand University of Medical Science, Birjand, Iran;

Telephone Number: 09155612135

Email Address: ghaderi@bums.ac.ir

following facial cosmetic procedures. In spite of the marked rise in nonsurgical cosmetic treatments, there is a deficiency of information about the effect of chemodervation and soft-tissue enhancement on psychosocial functioning (10).

On the other hand, due to cultural issues in some countries, only the faces of women can be exposed and the lips in the center of the face are always at the attention of the ladies who especially like their lips to be the same as a famous Hollywood Beautiful actress, Angelina Jolie, which I call the "Phenomenon of Angelinophilia" or "Ghaderi's Phenomenon". The attempt to have cosmetic facial surgeries has been increasing over the past years. Fillers for tissue enhancement are acquiring great popularity (11). For this purpose, the filler injection is used in the lip, especially the upper lip (12-15). Hyaluronic acid soft tissue fillers consist a range of productions (i.e., Voluma, Restylane Silk, Restylane, Restylane Lyft, Juvederm Ultra, Juvederm Ultra Plus, and Belotero Balance) that are applied frequently for the augmentation of facial features and facial rejuvenation. Although these productions are the same in many ways, they are not substitutable and have distinctive specifications that need to be noticed (13 -14).

There have been no large-scale analyses of complications related to filler injections (12). Liquid silicone which is also known as polydimethylsiloxane relates to a group of polymeric compounds that are frequently called silicones. These fillers have been applied as injectable filling agents for soft tissue enhancement and cosmetic lip augmentation (11). In a study, recently severe edema, erythema over cheeks and neck, and several irregularities were reported following lip augmentation with polydimethylsiloxane (11).

Despite the fact that specific adverse event profiles differ by material and injection site, common complications, such as swelling and infection are related to injectable filling agents. More severe complications include vascular events resulting in necrosis and visual loss; these complications are also increased in cases involving legal proceedings (12). Uncertified productions based on oleic solutions are used by uneducated personnel, thereby increasing the risk for specific adverse events, such as lipogranuloma formation, irritant and allergic contact dermatitis, and infections (15). On the other hand, the incidence of squamous cell carcinoma of the lip is high. It constitutes 25% to 30% of all oral cancers which is much more common in the lower lip than the upper lip (16- 19). The role of cumulative ultraviolet radiation in squamous cell carcinoma of

the lip is well established (19). These filler injections for cosmetic lip augmentation cause the upper lip mucosa to become more exposed to sun exposure

In my opinion, therefore, Ghaderi's phenomenon will greatly increase the risk of squamous cell carcinoma of the lip, especially in the upper lip. Accordingly, squamous cell carcinoma of the lip can be considered as a new complication following using filler injections for cosmetic lip augmentation.

"The future belongs to those who believe in the beauty of their dreams". Eleanor Roosevelt

Conflicts of Interest

There is no conflict of interest to be declared.

References

1. Giraldo-O'Meara M, Belloch A. Escalation from normal appearance related intrusive cognitions to clinical preoccupations in Body Dysmorphic Disorder: a cross-sectional study. *Psychiatry Res.* 2018; 265:137-43. [PMID: 29704773](#) [DOI: 10.1016/j.psychres.2018.04.047](#)
2. Hong K, Nezhgovorova V, Hollander E. New perspectives in the treatment of body dysmorphic disorder. *F1000Res.* 2018 ;7:361. [PMID: 29636904](#) [DOI: 10.12688/f1000research.13700.1](#)
3. Ghaderi R, Saadatjoo A. Quality of life in patients with hand eczema as health promotion: a case control study. *Acta Dermatovenerol Croat.* 2014; 22(1):32-9. [PMID: 24813839](#)
4. Ghaderi R, Saadatjoo A, Ghaderi F. Evaluating of life quality in patients with acne vulgaris using generic and specific questionnaires. *Dermatol Res Pract.* 2013; 2013:108624. [PMID: 24371434](#) [DOI: 10.1155/2013/108624](#)
5. Kamody RC, Thurston IB, Decker KM, Kaufman CC, Sonnevile KR, Richmond TK. Relating shape/weight based self-esteem, depression, and anxiety with weight and perceived physical health among young adults. *Body Image.* 2018; 25:168-76. [PMID: 29677688](#) [DOI: 10.1016/j.bodyim.2018.04.003](#)
6. Perdue TO, Schreier A, Swanson M, Neil J, Carels R. Evolving self view and body image concerns in female post-operative bariatric surgery patients. *J Clin Nurs.* 2018; 27(21-22):4018-27. [PMID: 29776012](#) [DOI: 10.1111/jocn.14527](#)
7. Scharschmidt D, Preiß S, Brähler E, Fischer T, Borkenhagen A. Body experience and self-esteem after minimally invasive skin rejuvenation: study of female patients using botulinum toxin A and/or dermal fillers. *Hautarzt.* 2017; 68(12):959-67. [PMID: 29052749](#) [DOI: 10.1007/s00105-017-4066-5](#)
8. Hashemi SAG, Edalatnoor B, Edalatnoor B, Niksun O.

- A comparison of body image concern in candidates for rhinoplasty and therapeutic surgery. *Electron Physician*. 2017; 9(9):5363-8. [PMID: 29038722](#) [DOI: 10.19082/5363](#)
9. Sobanko JF, Dai J, Gelfand JM, Sarwer DB, Percec I. Prospective cohort study investigating changes in body image, quality of life, and self-esteem following minimally invasive cosmetic procedures. *Dermatol Surg*. 2018; 44(8):1121-8. [PMID: 29659404](#) [DOI: 10.1097/DSS.0000000000001523](#)
 10. Imadojemu S, Sarwer DB, Percec I, Sonnad SS, Goldsack JE, Berman M, et al. Influence of surgical and minimally invasive facial cosmetic procedures on psychosocial outcomes: a systematic review. *JAMA Dermatol*. 2013; 149(11):1325-33. [PMID: 24068036](#) [DOI: 10.1001/jamadermatol.2013.6812](#)
 11. Abtahi-Naeini B, Faghihi G, Shahmoradi Z, Saffaei A. Filler migration and extensive lesions after lip augmentation: Adverse effects of polydimethylsiloxane filler. *J Cosmet Dermatol*. 2018; 17(6):996-9. [PMID: 29316184](#) [DOI: 10.1111/jocd.12476](#)
 12. Rayess HM, Svider PF, Hanba C, Patel VS, DeJoseph LM, Carron M, et al. A cross-sectional analysis of adverse events and litigation for injectable fillers. *JAMA Facial Plast Surg*. 2018; 20(3):207-214. [PMID: 29270603](#) [DOI: 10.1001/jamafacial.2017.1888](#)
 13. Gutowski KA. Hyaluronic acid fillers: science and clinical uses. *Clin Plast Surg*. 2016; 43(3):489-96. [PMID: 27363762](#) [DOI: 10.1016/j.cps.2016.03.016](#)
 14. Moradi A, Watson J. Current concepts in filler injection. *Facial Plast Surg Clin North Am*. 2015; 23(4):489-94. [PMID: 26505545](#) [DOI: 10.1016/j.fsc.2015.07.005](#)
 15. Kamouna B, Darlenski R, Kazandjieva J, Balabanova M, Dourmishev L, Negentsova Z, et al. Complications of injected vitamin E as a filler for lip augmentation: case series and therapeutic approach. *Dermatol Ther*. 2015; 28(2):94-7. [PMID: 25649466](#) [DOI: 10.1111/dth.12203](#)
 16. Liu F, Wang L, Pang S, Kan Q. Management of and risk factors for regional recurrence in upper lip squamous cell carcinoma. *Medicine (Baltimore)*. 2017; 96(45):e8270. [PMID: 29137010](#) [DOI: 10.1097/MD.00000000000008270](#)
 17. Assao A, Yoshino PM, Medeiros MCM, Carvalho AL, Soares FA, Kowalski LP, et al. Moesin involvement in oral carcinogenesis of the lower lip. *Anticancer Res*. 2018; 38(5):2755-60. [PMID: 29715096](#) [DOI: 10.21873/anticancer.12518](#)
 18. Han AY, Kuan EC, Mallen-St Clair J, Alonso JE, Arshi A, St John MA. Epidemiology of squamous cell carcinoma of the lip in the united states: a population-based cohort analysis. *JAMA Otolaryngol Head Neck Surg*. 2016; 142(12):1216-23. [PMID: 27832256](#) [DOI: 10.1001/jamaoto.2016.3455](#)
 19. Unsal AA, Unsal AB, Henn TE, Baredes S, Eloy JA. Cutaneous squamous cell carcinoma of the lip: A population-based analysis. *Laryngoscope*. 2018; 128(1):84-90. [PMID: 28573764](#) [DOI: 10.1002/lary.26704](#)